



JOURNAL OF BIODIVERSITY AND CONSERVATION

SHORT COMMUNICATION

Parasopubia delphiniifolia: a root parasitic unexplored plant

¹Sweta Mishra, ¹Sakti Kanta Rath and ²Sanjeet Kumar*¹Department of Life Sciences, Rama Devi Women's University, Bhubaneswar, Odisha, India²Ambika Prasad Research Foundation, Odisha, India

*Email-Id: sanjeetaprf@gmail.com

ARTICLE INFO

Article History

Received: 15 May 2023

Received in revised form: 19 June 2023

Accepted: 23 June 2023

Keywords: Parasitic plants, root parasite, *Parasopubia*, morphology

Abstract

Parasopubia delphiniifolia is an unexplored species belonging to the family Orobanchaceae. It is a root parasitic plant found near paddy fields, moist grasslands, and swampy areas. It has medicinal values. Still, it is unexplored. There is a need for more exploration work on it and its medicinal and ecological significance. Therefore, an attempt has been made to collect the information from field on *P. delphiniifolia*. In this communication, habit, habitat, plant parts, and uses are presented.

INTRODUCTION

Parasitic plants are a specialised plant group characterised by the ability to feed directly on other plants and penetrate either the shoots or roots of their host through a parasitic structure called haustoria (Wastewood et al. 2010; Sahu et al. 2018). Haustorium is a specialised organ composed of different tissue and cell types that might be homologous to the root or other subterranean plant structures, like tuber and rhizome, which penetrates host tissue like xylem

and phloem and creates a vascular bridge between parasite and host (Teixeira-Costa et al. 2020; Mishra et al. 2023). So, they are mainly of two types, i.e., stem parasites and root parasites (Sahu et al. 2018). In the case of root parasites, root hair-like structures are formed on the surface of the haustorium, which gradually elongates and captures the host vascular system, enabling the parasite to acquire water and nutrients and to modulate host physiology, probably via the secretion of virulence factors such as small molecules and proteins known as effectors (Saucet and Shirasu 2016; Sahu et al. 2018). Members of the family Orobanchaceae are mostly root holoparasites. They parasitize important crops and cause great losses in crop productivity. In this family, haustorium formation can occur at the meristematic tip of the parasite's primary root (called terminal haustorium) or at the transition zone on the side of a growing root (lateral haustorium) (Saucet and Shirasu 2016; Al-Mayah and Al-Asadi 2018). The genus *Parasopubia* Buch. -Ham. ex D. comprises about 30 species and shows some unique morphological characteristics. They live as hemiparasites in the dry or moist grasslands at higher altitudes or in or near swampy areas, including disturbed or cultivated ground such as rice paddies in the lowlands. These species are subshrubs or herbs with linear to lanceolate, simple, or dissected leaves. The androecium structure consists of four stamens, each with two different thecae, one normal-sized and fertile, the other sterile and reduced in size (Hofmann and Fischer 1998). Among these species, *P. delphiniifolia* (L.) H.-P. Hofm. & Eb. Fischer is an unexplored species with therapeutic values.

Parasopubia delphiniifolia

Parasopubia delphiniifolia (L.) H.-P. Hofm. & Eb. Fisch., Bot. Jahrb. Syst. 125(3): 357 2004. (syn: *Euphrasia coromandelina* Rottl. ex Spreng.; *Gerardia delphinifolia* L.; *Gerardia heyneana* Benth.; *Lophanthera delphinifolia* (L.) Raf.; *Sopubia delphinifolia* (L.) G.Don).

MORPHOLOGY

Perennial, erect herb, stem hispid, about 40-60 cm tall. Leaves pinnatifid below, upper ones opposite, linear, filiform, 2-3.5 cm long, margin entire, apex acute. Flowers axillary, solitary, become racemose above, pedicellate, calyx tube 10- ribbed, glabrous, corolla purple with violet streaks, 8-16 mm long; stamens 4. Capsule oblong-ellipsoid, 5 x 3.5 mm, seeds oblong, cuneate (Plate 1).

Flowering & Fruiting: August- December



Plate 1: Habit (a), flower (b) and habitat of *P. delphiniifolia*

Habitat: Along the borders of paddy fields, roadsides, in moist grasslands or, near swampy areas

Distribution: India, Sri Lanka

Medicinal uses: Fresh leaf extract of *P. delphiniifolia* is externally applied on cuts and wounds. Its juice is reported to be an abortifacient.

FUTURE ASPECTS

The present study could provide baseline data for further experimental work on unexplored parasitic plant species of Odisha. Field surveys and correct identification will help in a new sight into the physiology, medicinal uses, ecological and evolutionary biology of parasitic plants.

REFERENCES

- Al-Mayah AA and Al-Asadi WM. (2018). An Account of *Orobanche* L. (Orobanchaceae) in Iraq. Biological and Applied Environmental Research. 2(1): 22-33.
- Hofmann HP and Fischer E. (1998). Preliminary revision of the genus *Sopubia* (Scrophulariaceae) in Madagascar. Adansonia. 20(2): 299-312.
- Mishra S, Rath SK, Jeet K, Woldeamanuel MM and Kumar S. (2023). *Cassytha filiformis*: a parasitic climber of Odisha, India. In Medicinal Climber. 1-41.
- Sahu SC, Singh NR and Kumar S. (2018). Parasitic plants of Odisha. APRF Publisher, Odisha, India.
- Saucet SB and Shirasu K. (2016). Molecular Parasitic Plant–Host Interactions. PLoS Pathogens. 12:(12). e1005978. DOI: 10.1371/journal.ppat.1005978.
- Teixeira-Costa L, Ocampo G and Ceccantini G. (2020). Morphogenesis and evolution of mistletoes' haustoria. In Plant Ontogeny. 107-157.
- Westwood JH, Yoder JI, Timko MP and Depamphilis CW. (2010). The evolution of parasitism in plants. Trends in Plant Sciences. 15: 227–235. DOI: 10.1016/j.tplants.2010.01.004.